

Correction

Intraocular pressure in genetically distinct mice: an update and strain survey: Correction

Olga V Savinova¹, Fumihiko Sugiyama², Janice E Martin^{1,3},
Stanislav I Tomarev⁴, Beverly J Paigen¹, Richard S Smith^{1,3} and
Simon WM John^{*1,3,5}

Address: ¹The Jackson Laboratory, 600 Main Street, Bar Harbor, ME, 04609, USA, ²Laboratory Animal Resource Center, University of Tsukuba, Tsukuba, 305, Japan, ³Howard Hughes Medical Institute at The Jackson Laboratory, 600 Main Street, Bar Harbor, ME, 04609, USA, ⁴Laboratory of Molecular and Developmental Biology, National Eye Institute, NIH, Bethesda, MD, 20892-3655, USA and ⁵Department of Ophthalmology, Tufts University School of Medicine, Boston, MA, 02155, USA

E-mail: Olga V Savinova - ovs@jax.org; Fumihiko Sugiyama - bunbun@sakura.cc.tsukuba.ac.jp; Janice E Martin - jemartin@jax.org; Stanislav I Tomarev - tomarev@intra.nei.nih.gov; Beverly J Paigen - bip@jax.org; Richard S Smith - rss@jax.org; Simon WM John* - swmj@jax.org

*Corresponding author

Published: 26 September 2001

Received: 26 September 2001

BMC Genetics 2001, 2:16

Accepted: 26 September 2001

This article is available from: <http://www.biomedcentral.com/1471-2156/2/16>

© 2001 Savinova et al; licensee BioMed Central Ltd. Verbatim copying and redistribution of this article are permitted in any medium for any non-commercial purpose, provided this notice is preserved along with the article's original URL. For commercial use, contact info@biomedcentral.com

Editorial note

The original article can be viewed at [<http://www.biomedcentral.com/1471-2156/2/12>] [1].

Correction

1) Legend to figure 7 should read: "There is no strong correlation between the average systolic blood pressure of each strain and the average IOP of that strain ($R^2 = 0.1$). All mice were female and 2 to 4 months old, with the exception that the female CBA/CaJ used for IOP assessment were 6 to 7 months old. If the CBA/CaJ mice are excluded, R^2 drops to 0.01. Of ARK/J, C57BL/6J and SJL/J, strain ARK/J had the highest IOP and strain SJL/J the lowest IOP.

2) In table 1, the strain NZB/BiNJ should read: NZB/B1NJ.

3) In the results section dealing with *Myoc* alleles, the MYOC and Myoc should both be in italics.

References

1. Savinova OV, Sugiyama F, Martin JE, Tomarev SI, Paigen BJ, Smith RS, John SWM: **Intraocular pressure in genetically distinct mice: an update and strain survey.** *BMC Genetics* 2001, 2:12